



# High concordance of 70-gene recurrence risk signature and 80-gene molecular subtyping signature between core needle biopsy and surgical resection specimens in early-stage breast cancer

Jennifer A. Crozier MD<sup>1</sup> | Julie Barone DO<sup>2</sup> | Pat Whitworth MD<sup>3</sup> |  
Abraham Cheong MD<sup>4</sup> | Robert Maganini MD<sup>5</sup> | Jose Perez Tamayo MD<sup>6</sup> |  
Patricia Dauer PhD<sup>7</sup>  | Shiyu Wang MS<sup>7</sup> | William Audeh MD<sup>7</sup> |  
Annuska M. Glas PhD<sup>8</sup> 

<sup>1</sup>Division of Hematology & Oncology, Baptist MD Anderson, Jacksonville, Florida, USA

<sup>2</sup>SCL Health, St. Joseph's Hospital, Denver, Colorado, USA

<sup>3</sup>Department of Surgery, Nashville Breast Center, Nashville, Tennessee, USA

<sup>4</sup>Division of Hematology & Oncology, Southeast Georgia Health System, Brunswick, Georgia, USA

<sup>5</sup>Division of Oncology, AMITA Health Alexian Brothers, Elk Grove Village, Illinois, USA

<sup>6</sup>Department of Radiology, Ogden Regional Medical Center, Ogden, Utah, USA

<sup>7</sup>Division of Medical Affairs, Agendia Inc., Irvine, California, USA

<sup>8</sup>Division of R&D, Agendia NV, Amsterdam, The Netherlands

## Correspondence

Annuska M. Glas, PhD, Division of R&D and Innovation, Agendia NV, Radarweg 60, 1043 NT Amsterdam, The Netherlands.  
Email: [annuska.glas@agendia.com](mailto:annuska.glas@agendia.com)

## Abstract

**Background and Objectives:** With increased neoadjuvant therapy recommendations for early-stage breast cancer patients due to the COVID-19 pandemic, it is imperative that molecular diagnostic assays provide reliable results from preoperative core needle biopsies (CNB). The study objective was to determine the concordance of MammaPrint and Blueprint results between matched CNB and surgical resection (SR) specimens.

**Methods:** Matched tumor specimens ( $n = 121$ ) were prospectively collected from women enrolled in the FLEX trial (NCT03053193). Concordance is reported using overall percentage agreement and Cohen's kappa coefficient. Correlation is reported using Pearson correlation coefficient.

**Results:** We found good concordance for MammaPrint results between matched tumor samples (90.9%,  $\kappa = 0.817$ ), and a very strong correlation of MammaPrint indices ( $r = 0.94$ ). The concordance of Blueprint subtyping in matched samples was also excellent (98.3%).

**Conclusions:** CNB samples demonstrated high concordance with paired SR samples for MammaPrint risk classification and Blueprint molecular subtyping, suggesting that physicians are provided with accurate prognostic information that can be used to guide therapy decisions.

## KEYWORDS

Blueprint, breast cancer, core needle biopsy, COVID-19 pandemic, genomic profile, MammaPrint

**Abbreviations:** CNB, core needle biopsy; EBC, early-stage breast cancer; ER, estrogen receptor; FFPE, formaldehyde-fixed paraffin-embedded; HER2, human epidermal growth factor receptor 2; HR, hormone receptor; IHC, immunohistochemistry; NPV, negative predictive value; PPV, positive predictive value; PR, progesterone receptor; SR, surgical resection; TN, triple negative.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2021 The Authors. *Journal of Surgical Oncology* published by Wiley Periodicals LLC

## 1 | INTRODUCTION

Neoadjuvant systemic therapy has been increasingly used as clinical trial results have demonstrated no difference in disease-free survival or overall survival compared with patients who received adjuvant therapy.<sup>1-3</sup> In light of the extraordinary circumstances that the COVID-19 pandemic presented last year, use of neoadjuvant therapy has accelerated further. Health experts and professional societies, including the American College of Surgeons, published treatment guidelines to conserve supplies, triage cases, and reduce risk without compromising patient care.<sup>4-7</sup> As a result, neoadjuvant therapy became even more prevalent in breast cancer treatment plans which requires prognostic information to be obtained from a core needle biopsy (CNB) before preoperative therapy.

Multigene assays, like MammaPrint, provide prognostic information that can guide physicians on therapy decisions for patients. However, multigene assays have typically been developed and tested using primary surgical specimens in an adjuvant setting. Only a few studies have determined the performance of multigene assays using CNB in recent years.<sup>8-11</sup> In a large unpaired study, Jakubowski et al. found a similar range of Oncotype Dx Recurrence Scores between CNB and surgical resection (SR) samples (10-22 vs. 11-22, respectively).<sup>8</sup> In three smaller studies using matched CNB and SR samples, researchers reported Pearson correlation coefficients ranging from 0.20 to 0.99 and overall concordance ranging from 72.0% to 95.0% using either EndoPredict, GenesWell, or Oncotype Dx multigene assays.<sup>9-11</sup>

MammaPrint is a 70-gene assay that can be used to predict the likelihood of recurrence and response to chemotherapy.<sup>12,13</sup> Patients identified as MammaPrint Low Risk can safely forego chemotherapy without compromising outcome, compared to those identified as High Risk for whom chemotherapy is recommended.<sup>13</sup> MammaPrint has demonstrated precision (99.0%) and reproducibility (98.9%) in fresh frozen tissue, with 95% agreement between two sample sites from the same tumor.<sup>14</sup> Use of formalin-fixed paraffin-embedded (FFPE) tissue, which make up most CNB samples, demonstrated very strong correlation ( $r = 0.92$ ) and a 91.5% concordance of MammaPrint classification results with fresh tissue.<sup>15</sup>

In addition to risk classification, multigene assays, such as BluePrint, can be utilized to reliably determine the molecular subtype of women with early-stage breast cancer (EBC).<sup>16</sup> We have previously tested the performance of BluePrint molecular subtyping in comparison with immunohistochemistry (IHC) using multiple cohorts and have demonstrated overall subtype reclassification of up to 30%.<sup>17-19</sup> Additionally, tumors reclassified by MammaPrint and BluePrint exhibited more accurate pathological complete response (pCR) rates compared to pCR rates based on their respective clinical subtype.<sup>16,18</sup> These results support the importance of multigene assays in treatment decisions.

Though CNB samples have been used for MammaPrint and BluePrint in development and validation studies, as well as in clinical trials I-SPY2 (NCT01042379) and NBRST (NCT01479101), no study has directly compared CNB and SR specimens. In our current study,

we have prospectively collected 139 matched CNB and SR specimens from women with confirmed EBC enrolled in the ongoing FLEX study (NCT03053193). The objective of our study is to determine the concordance of MammaPrint and BluePrint results between CNB and SR, to ensure reliable prognostic information can be captured from a CNB.

## 2 | MATERIALS AND METHODS

### 2.1 | Patient cohort

FLEX is an ongoing, multiinstitutional prospective study of patients with Stage I-III EBC. This study was conducted in accordance with the ethical standards established by the Declaration of Helsinki. The protocol was approved by Institutional Review Boards at all participating sites and registered with ClinicalTrials.gov (NCT03053193). Enrolled patients receive a MammaPrint test with or without BluePrint molecular subtyping and consent to clinically annotated gene expression data collection. Patients within the FLEX study ( $n = 139$ ) with matched CNB and SR specimens were prospectively collected from six institutions. We excluded 18 patients due to neoadjuvant treatment, which resulted in 121 eligible matched CNB and SR tissues for this study.

### 2.2 | Molecular classification

For this study, patients were identified in our FLEX patient database by their CNB MammaPrint result, and corresponding SR tissue was requested from their respective local institution. Upon receipt of SR FFPE tissue blocks at Agendia (Irvine, CA), sections were prepared. In accordance with diagnostic quality controls and standards, one section per sample was reviewed internally to verify >30% tumor cellularity. RNA was isolated with the RNeasy FFPE kit (Qiagen), and concentration measured by NanoDrop 2000 (Thermo Fisher Scientific). Complementary DNA was labeled and hybridized to 44k arrays and scanned using a dual laser scanner (Agilent Technologies) as previously described.<sup>15,20,21</sup>

MammaPrint Low Risk tumors have a MammaPrint index of >0.000; High Risk tumors have a MammaPrint Index of  $\leq 0.000$ . BluePrint classifies tumors as Luminal-type, Basal-type, or human epidermal growth factor receptor 2 (HER2)-type. Each tumor sample is scored for all three subtypes, with the highest index representing the respective categorical tumor subtype.<sup>20</sup> Together, MammaPrint and BluePrint stratify Luminal-type tumors into Luminal A (MammaPrint Low Risk) or Luminal B (MammaPrint High Risk). For this study, borderline samples (MammaPrint index between -0.05 and 0.05) were excluded due to a higher probability of technical inaccuracy, as reported in the FDA 510k summary (#K070675). It should be noted that borderline results occur in less than 5% of the total cases, and the likelihood of a borderline result is independent of sample type (i.e., CNB or SR).

## 2.3 | Statistical analysis

A power analysis was conducted using Lin's Concordance Correlation Coefficient to evaluate the reproducibility of the two methods. The two one-sided option was chosen to test equivalence, assuming equivalence limit range of 90.0%–99.9% and expected correlation to be 0.90; 121 samples provide a power of 93.2%. Descriptive statistics were used to summarize patient clinicopathological characteristics. The primary objective of this study was to evaluate the concordance of MammaPrint results between CNB and SR, measured using overall percentage agreement, positive predictive value (PPV, High Risk), negative predictive value (NPV, Low Risk), and Cohen's kappa coefficient ( $k$ ). The secondary objective was to determine the concordance of Blueprint molecular subtypes between matched CNB and SR samples. Pearson correlation coefficient ( $r$ ) and Pearson correlation test were calculated using the MammaPrint index or Blueprint indices for CNB and SR specimens. Pearson Correlation values were interpreted as outlined by Schober, et. al.,<sup>22</sup> where  $r = 0.40$ – $0.69$  is considered a “moderate correlation”,  $r = 0.70$ – $0.89$  is a “strong correlation”, and  $0.90$ – $1.00$  is a “very strong correlation”. A  $p$  value less than 0.05 is considered statistically significant. Statistical analyses were performed using GraphPad Prism (version 9.0.2) and R (version 4.0.5).

## 3 | RESULTS

### 3.1 | Patient characteristics

A total of 121 patients from the FLEX study database with diagnostic MammaPrint and Blueprint results with matched CNB and SR specimens were included in this study. Table 1 summarizes clinicopathological characteristics of this cohort. Out of the 119 patients with documented age, the majority of patients were over the age of 50 (97/119; 81.5%). Of the 114 patients with clinical subtyping data, most patients (97/114; 85.1%) had hormone receptor (HR) positive/HER2 negative tumors, 10 (8.8%) had HR positive/HER2 positive tumors, and 7 (6.1%) had triple negative (TN) tumors. No patients in this cohort had HR negative/HER2 positive tumors. Out of 119 patient tumors, 36 (30.3%) were low grade (G1), 53 (44.5%) were intermediate grade (G2), and 30 (25.2%) were high grade (G3).

### 3.2 | Concordance of MammaPrint results between CNB and SR tumor specimens

Overall, 50 patients had High Risk CNB and SR specimens and 60 had Low Risk CNB and SR specimens, resulting in 90.9% overall agreement ( $\kappa = 0.817$ ), 95.2% NPV, and 86.2% PPV (Table 2). Out of the discordant samples, eight were High Risk on CNB and Low Risk on SR, whereas three were Low Risk on CNB and High Risk on SR. A Pearson correlation test of MammaPrint indices between 121 CNB and SR specimens was performed and resulted in a very strong correlation of  $r = 0.94$  ( $p < 0.0001$ ) (Figure 1).

**TABLE 1** Patient tumor clinical characteristics

	Patient number (%) Total $n = 121$
Age <sup>a</sup>	
>50	97 (81.5%)
≤50	22 (18.5%)
Clinical subtype <sup>b*</sup>	
HR positive/HER2 negative	97 (85.1%)
HR positive/HER2 positive	10 (8.8%)
Triple negative	7 (6.1%)
Tumor grade <sup>c</sup>	
G1 low grade	36 (30.3%)
G2 intermediate grade	53 (44.5%)
G3 high grade	30 (25.2%)
Tumor stage <sup>d</sup>	
T1	76 (66.7%)
T2	33 (28.9%)
T3	5 (4.4%)
Nodal status <sup>e</sup>	
Negative	53 (76.8%)
Positive	16 (23.2%)
Surgery type <sup>f</sup>	
Lumpectomy	30 (62.5%)
Mastectomy	18 (37.5%)
Method of detection <sup>g</sup>	
Clinical palpation/finding	3 (2.5%)
Screening mammogram	90 (75.6%)
Self-exam/patient discovered	26 (21.8%)


Note: <sup>a,c,g</sup> $n = 119$ , 1.7% unknown clinical data; <sup>b,d</sup> $n = 114$ , 5.8% unknown clinical data; \*HER2 equivocal patients ( $n = 3$ ) counted as HER2 negative; <sup>e</sup> $n = 69$ , 41.3% unknown clinical data; <sup>f</sup> $n = 48$ , 60.3% unknown clinical data.

Abbreviation: HR, hormone receptor.

**TABLE 2** Concordance of MammaPrint results between CNB and SR

MammaPrint result	SR		Total
	High risk	Low risk	
CNB			
High risk	50	8	58
Low risk	3	60	63
Total	53	68	121

Abbreviations: CNB, core needle biopsy; SR, surgical resection.



**FIGURE 1** Correlation of MammaPrint Index between core needle biopsy (CNB) and surgical resection (SR). MammaPrint index was determined for each matched CNB and SR tumor specimen ( $n = 121$ ) and the correlation of matched samples was determined using Pearson correlation coefficient

**TABLE 3** Concordance of Blueprint subtyping classification between CNB and SR

Blueprint result	SR			Total
	Luminal-type	HER2-type	Basal-type	
CNB				
Luminal-type	105	0	1	106
HER2-type	0	2	0	2
Basal-type	1	0	12	13
Total	106	2	13	121

Abbreviations: CNB, core needle biopsy; SR, surgical resection.

### 3.3 | Concordance of Blueprint molecular subtyping between CNB and SR tumor specimens


CNB and SR tumors were in agreement for 105/106 Luminal-type (99.1%), 2/2 HER2-type (100%), and 12/13 Basal-type (92.3%) tumors (Table 3). Overall, we determined the concordance of Blueprint between CNB and SR to be 98.3%. In addition, we observed very strong correlations of Blueprint index scores between CNB and SR specimens for Luminal-type ( $r = 0.92$ ;  $p < 0.0001$ ) and Basal-type ( $r = 0.97$ ;  $p < 0.0001$ ) tumors (Figure 2). We did not determine the Pearson correlation coefficient for HER2-type patients due to too few by Blueprint ( $n = 2$ ).

## 4 | DISCUSSION

In recent years, neoadjuvant therapy has been increasingly used for tumor and/or nodal downstaging, monitoring treatment response, and to allow time for genomic testing, which has expanded the number of patients receiving genomic testing and helped streamline care.<sup>23</sup> With an overall increase in neoadjuvant therapy and where triaging patients for neoadjuvant therapy is recommended (e.g., during the COVID-19 pandemic<sup>4–7</sup>), physicians can use molecular subtyping and risk classification information in addition to clinical pathology on preoperative core biopsies, which can guide therapy decision making. The standard of care method for breast cancer subtype classification and treatment decisions are based on clinical features (e.g., tumor grade, lymph node status, receptor status, and Ki-67) as determined by IHC/FISH. Multiple studies have analyzed the concordance of CNB and SR specimens in tumor grade, with overall agreement ranging from 64% to 77%.<sup>24–27</sup> In addition, numerous reports comparing SR and CNB samples in estrogen receptor, progesterone receptor, HER2, and Ki-67 IHC staining have been performed with overall agreement ranging from 84.0% to 99.1%, 77.9% to 94.3%, 80.0% to 98.8%, and 79.5% to 87.0% respectively, and a wide range of overall concordance (75%–97%).<sup>26,28–35</sup>

Based on our previous performance and concordance studies using MammaPrint and Blueprint,<sup>20,36,37</sup> we anticipated the overall agreement of MammaPrint results between CNB and SR tumor samples to be similar. As expected, we observed an overall agreement of 90.9%. In the current study, 11/121 samples were discordant between CNB and SR, with 8 samples having High Risk CNB results and Low Risk SR results, and 3 samples with Low Risk CNB results and High Risk SR results. It is important to note that discordance between sample type reflects the complex tumor heterogeneity, where sample size and sample location within the tumor can lead to differences in risk assessment, but both are accurate results. Therefore, High Risk CNB results reflect a portion of the tumor that was High Risk, even if the SR specimen was Low Risk, and recommended therapeutic options would follow guidelines for High Risk tumors. Importantly, clinically relevant discordant cases that affect treatment decisions are cases in which CNB is Low Risk and SR is High Risk, which account for approximately 2.5% of patients in the current analysis. In these few cases, the High Risk tumor biology not captured on the preoperative CNB would potentially underestimate treatment decisions. Overall, these results confirm that MammaPrint risk classification from CNB are in high agreement with SR and provide consistent and accurate results. Thus, for more than 97% of patients in this study, treatment decisions and potential outcome are precise and consistent based on MammaPrint testing of the CNB.

In addition to our primary analysis, we determined the concordance of Blueprint genomic subtyping between the matched samples. Out of 121 tumors, 119 were in agreement, resulting in an overall concordance of 98.3%; superior to concordance rates of



**FIGURE 2** Correlation of Blueprint Index between CNB and SR. Blueprint generates three indices corresponding to Luminal-type, HER2-type, and Basal-type. Indices for (A) Luminal-type, and (B) Basal-type were graphed and a Pearson correlation test was performed for each. (C) All BluePrint indices ( $n = 121$ ) with Luminal-type in blue, HER2-type in green, and Basal-type in red. CNB, core needle biopsy; SR, surgical resection

75.0%–87.5% among intrinsic biological subtypes based on IHC assessment.<sup>26–28</sup> Molecular subtyping results have demonstrated more accurate prediction of a pCR and outcome in comparison with IHC assessment.<sup>17,38,39</sup> Although reclassification by BluePrint was not an objective of this study, it was notable that 8 of the 10 patients clinically identified as HER2+ were reclassified as Luminal-type by BluePrint. Additionally, a CNB provides accurate and fast results that can be used for treatment decisions, and shortens time-to-treat, which can ultimately improve patient outcomes.<sup>40</sup> A limitation of our paired study is data maturity, in which patient follow-up data is currently unavailable to correlate outcome with MammaPrint and BluePrint results from CNB and SR samples. However, several studies, including the NBRST trial (NCT01479101), ISPY1 (NCT00033397), and ISPY2 (NCT01042379), have demonstrated accurate prediction in neoadjuvant treatment response and long-term outcome by MammaPrint and BluePrint on core needle biopsies.<sup>17,39,41–43</sup>

## 5 | CONCLUSIONS

In summary, this analysis represents the largest powered study using prospective real-world data to evaluate the concordance of a genomic assay on matched CNB and SR samples. The high concordance rates of MammaPrint and BluePrint results between paired samples strongly support the utility of these assays to obtain reliable prognostic information on core biopsy tissue, which can guide timely and appropriate treatment decisions.

## ACKNOWLEDGMENTS

We would like to thank all the patients who volunteered to participate in FLEX, as well as the FLEX site coordinators and investigators. We would also like to thank the following people for their contribution to this manuscript: Jeffrey Falk, Jia-Perng Wei, Sammy Mee, Jake Ruby, Suoyi Yang, Yen Huynh, Anke Witteveen, Christine Finn, Kate Corcoran, Christa Dreezen, Andrea Menicucci, Annie Tran, Erin

Yoder, and Bastiaan van der Baan. Research was supported by Agendia Inc.

### CONFLICT OF INTERESTS

Jennifer A. Crozier, Julie Barone, Pat Whitworth, Abraham Cheong, Robert Maganini, and Jose Perez Tamayo are FLEX principal investigators and have contracted research with Agendia Inc. Jennifer A. Crozier and Robert Maganini received honoraria as part of Speaker's Bureau for Agendia Inc. Patricia Dauer, Shiyu Wang, and William Audeh, are noncommercial employees of Agendia Inc, Irvine, CA. Annuska Glas is a noncommercial employee of Agendia NV, Amsterdam, The Netherlands. Annuska Glas is a coinventor of the BluePrint 80-gene signature and is a full-time employee of Agendia, NV (Patent Numbers: 9175351, 10072301). No other conflict of interests were reported.

### AUTHOR CONTRIBUTIONS

**Patricia Dauer, Shiyu Wang, William Audeh, and Annuska Glas:** contributed to data analysis and interpretation. **Jennifer A. Crozier, Patricia Dauer, and Shiyu Wang:** contributed to manuscript preparation. All authors participated in the review and editing of the manuscript for publication. All authors contributed to the study design and conceptualization. All authors contributed to collection and assembly of data.

### DATA AVAILABILITY STATEMENT

The clinical datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request. Raw array data have not been made publicly available as part of a collaboration agreement with the diagnostic company Agendia Inc.

### ORCID

Patricia Dauer  <http://orcid.org/0000-0002-8374-5512>

Annuska M. Glas  <https://orcid.org/0000-0002-0775-138X>

### REFERENCES

1. Wolmark N, Wang J, Mamounas E, Bryant J, Fisher B. Preoperative chemotherapy in patients with operable breast cancer: nine-year results from National Surgical Adjuvant Breast and Bowel Project B-18. *J Natl Cancer Inst Monogr*. 2001;30:96-102.
2. van der Hage JA, van de Velde CJ, Julien JP, Tubiana-Hulin M, Vandervelden C, Duchateau L. Preoperative chemotherapy in primary operable breast cancer: results from the European Organization for Research and Treatment of Cancer trial 10902. *J Clin Oncol*. 2001;19(22):4224-4237.
3. Mauri D, Pavlidis N, Ioannidis JP. Neoadjuvant versus adjuvant systemic treatment in breast cancer: a meta-analysis. *J Natl Cancer Inst*. 2005;97(3):188-194.
4. Ueda M, Martins R, Hendrie PC, et al. Managing cancer care during the COVID-19 pandemic: agility and collaboration toward a common goal. *J Natl Compr Canc Netw*. 2020;18(4):1-4.
5. Curigliano G, Cardoso MJ, Poortmans P, et al. Recommendations for triage, prioritization and treatment of breast cancer patients during the COVID-19 pandemic. *Breast*. 2020;52:8-16.
6. Surgeons ACo. COVID 19: Elective Case Triage Guidelines for Surgical Care. 2020.
7. Oncology SoS. Resource for Management Options of Breast Cancer During COVID-19. 2020.
8. Jakubowski DM, Bailey H, Abran J, et al. Molecular characterization of breast cancer needle core biopsy specimens by the 21-gene breast recurrence score test. *J Surg Oncol*. 2020;122(4):611-618.
9. Müller BM, Brase JC, Haufe F, et al. Comparison of the RNA-based EndoPredict multigene test between core biopsies and corresponding surgical breast cancer sections. *J Clin Pathol*. 2012;65(7):660-662.
10. Lee J, Lee EH, Park HY, et al. Efficacy of an RNA-based multigene assay with core needle biopsy samples for risk evaluation in hormone-positive early breast cancer. *BMC Cancer*. 2019;19(1):388.
11. Qi P, Yang Y, Bai QM, et al. Concordance of the 21-gene assay between core needle biopsy and resection specimens in early breast cancer patients. *Breast Cancer Res Treat*. 2021;186(2):327-342.
12. van 't Veer LJ, Veer, Dai H, van de Vijver MJ, et al. Gene expression profiling predicts clinical outcome of breast cancer. *Nature*. 2002;415:530-536.
13. Cardoso F, van't Veer LJ, Bogaerts J, et al. 70-Gene signature as an aid to treatment decisions in early-stage breast cancer. *N Engl J Med*. 2016;375(8):717-729.
14. Delahaye LJ, Wehkamp D, Floore AN, Bernards R, Van 't Veer L, Glas AM. Performance characteristics of the MammaPrint breast cancer diagnostic gene signature. *Personalized Medicine*. 2013;10(8):10-811.
15. Sapino A, Roepman P, Linn SC, et al. MammaPrint molecular diagnostics on formalin-fixed, paraffin-embedded tissue. *J Mol Diagn*. 2014;16(2):190-197.
16. Krijgsman O, Roepman P, Zwart W, et al. A diagnostic gene profile for molecular subtyping of breast cancer associated with treatment response. *Breast Cancer Res Treat*. 2012;133(1):37-47.
17. Whitworth P, Stork-Sloots L, de Snoo FA, et al. Chemosensitivity predicted by BluePrint 80-gene functional subtype and MammaPrint in the Prospective Neoadjuvant Breast Registry Symphony Trial (NBRST). *Ann Surg Oncol*. 2014;21(10):3261-3267.
18. Viale G, de Snoo FA, Slaets L, et al. Immunohistochemical versus molecular (BluePrint and MammaPrint) subtyping of breast carcinoma. Outcome results from the EORTC 10041/BIG 3-04 MIND-ACT trial. *Breast Cancer Res Treat*. 2018;167(1):123-131.
19. Nguyen B, Cusumano PG, Deck K, et al. Comparison of molecular subtyping with BluePrint, MammaPrint, and TargetPrint to local clinical subtyping in breast cancer patients. *Ann Surg Oncol*. 2012;19(10):3257-3263.
20. Mitterpergher L, Delahaye LJ, Witteveen AT, et al. Performance characteristics of the BluePrint(R) breast cancer diagnostic test. *Transl Oncol*. 2020;13(4):100756.
21. Glas AM, Floore A, Delahaye LJ, et al. Converting a breast cancer microarray signature into a high-throughput diagnostic test. *BMC Genomics*. 2006;7:278.
22. Schober P, Boer C, Schwarte LA. Correlation coefficients: appropriate use and interpretation. *Anesth Analg*. 2018;126(5):1763-1768.
23. Cain H, Macpherson IR, Beresford M, Pinder SE, Pong J, Dixon JM. Neoadjuvant therapy in early breast cancer: treatment considerations and common debates in practice. *Clin Oncol (R Coll Radiol)*. 2017;29(10):642-652.
24. Connor CS, Tawfik OW, Joyce AJ, Davis MK, Mayo MS, Jewell WR. A comparison of prognostic tumor markers obtained on image-guided breast biopsies and final surgical specimens. *Am J Surg*. 2002;184(4):322-324.
25. Harris GC, Denley HE, Pinder SE, et al. Correlation of histologic prognostic factors in core biopsies and therapeutic excisions of invasive breast carcinoma. *Am J Surg Pathol*. 2003;27(1):4-5.

26. Burge CN, Chang HR, Apple SK. Do the histologic features and results of breast cancer biomarker studies differ between core biopsy and surgical excision specimens? *Breast*. 2006;15(2):167-172.
27. Daveau C, Baulies S, Lalloum M, et al. Histological grade concordance between diagnostic core biopsy and corresponding surgical specimen in HR-positive/HER2-negative breast carcinoma. *Br J Cancer*. 2014;110(9):2195-2200.
28. You K, Park S, Ryu JM, et al. Comparison of core needle biopsy and surgical specimens in determining intrinsic biological subtypes of breast cancer with immunohistochemistry. *J Breast Cancer*. 2017;20(3):297-303.
29. Lorgis V, Algros MP, Villanueva C, et al. Discordance in early breast cancer for tumour grade, estrogen receptor, progesterone receptors and human epidermal receptor-2 status between core needle biopsy and surgical excisional primary tumour. *Breast*. 2011;20(3):284-287.
30. Dekker TJ, Smit VT, Hooijer GK, et al. Reliability of core needle biopsy for determining ER and HER2 status in breast cancer. *Ann Oncol*. 2013;24(4):931-937.
31. Tamaki K, Sasano H, Ishida T, et al. Comparison of core needle biopsy (CNB) and surgical specimens for accurate preoperative evaluation of ER, PgR and HER2 status of breast cancer patients. *Cancer Sci*. 2010;101(9):2074-2079.
32. Arnedos M, Nerurkar A, Osin P, A'Hern R, Smith IE, Dowsett M. Discordance between core needle biopsy (CNB) and excisional biopsy (EB) for estrogen receptor (ER), progesterone receptor (PgR) and HER2 status in early breast cancer (EBC). *Ann Oncol*. 2009;20(12):1948-1952.
33. Mann GB, Fahey VD, Feleppa F, Buchanan MR. Reliance on hormone receptor assays of surgical specimens may compromise outcome in patients with breast cancer. *J Clin Oncol*. 2005;23(22):5148-5154.
34. Chen X, Sun L, Mao Y, et al. Preoperative core needle biopsy is accurate in determining molecular subtypes in invasive breast cancer. *BMC Cancer*. 2013;13:390.
35. Kombak FE, Şahin H, Mollamemişoğlu H, et al. Concordance of immunohistochemistry between core needle biopsy and surgical resection of breast cancer. *Turk J Med Sci*. 2017;47(6):1791-1796.
36. Mitterpergher L, de Ronde JJ, Nieuwland M, et al. Gene expression profiles from formalin fixed paraffin embedded breast cancer tissue are largely comparable to fresh frozen matched tissue. *PLoS One*. 2011;6(2):e17163.
37. Beumer I, Witteveen A, Delahaye L, et al. Equivalence of MammaPrint array types in clinical trials and diagnostics. *Breast Cancer Res Treat*. 2016;156(2):279-287.
38. Whitworth P, Beitsch P, Mislowsky A, et al. Chemosensitivity and endocrine sensitivity in clinical luminal breast cancer patients in the prospective neoadjuvant breast registry symphony trial (NBRST) predicted by molecular subtyping. *Ann Surg Oncol*. 2017;24(3):669-675.
39. Whitworth P, Pellicane JV, Baron P, et al. Abstract PD9-01: 5-year outcomes in the NBRST trial: Preoperative MammaPrint and BluePrint breast cancer subtype is associated with neoadjuvant treatment response and survival. *Cancer Res*. 2021;81(4 Suppl):PD9-01-PD09-01.
40. Bleicher RJ. Timing and delays in breast cancer evaluation and treatment. *Ann Surg Oncol*. 2018;25(10):2829-2838.
41. Gluck S, de Snoo F, Peeters J, Stork-Sloots L, Somlo G. Molecular subtyping of early-stage breast cancer identifies a group of patients who do not benefit from neoadjuvant chemotherapy. *Breast Cancer Res Treat*. 2013;139(3):759-767.
42. Esserman LJ, Berry DA, Cheang MC, et al. Chemotherapy response and recurrence-free survival in neoadjuvant breast cancer depends on biomarker profiles: results from the I-SPY 1 TRIAL (CALGB 150007/150012; ACRIN 6657). *Breast Cancer Res Treat*. 2012;132(3):1049-1062.
43. Wolf DM, Yau C, Sanil A, et al. DNA repair deficiency biomarkers and the 70-gene ultra-high risk signature as predictors of veliparib/carboplatin response in the I-SPY 2 breast cancer trial. *NPJ Breast Cancer*. 2017;3:31.

**How to cite this article:** Crozier JA, Barone J, Whitworth P, et al. High concordance of 70-gene recurrence risk signature and 80-gene molecular subtyping signature between core needle biopsy and surgical resection specimens in early-stage breast cancer. *J Surg Oncol*. 2022;125:596-602.  
doi:10.1002/jso.26780